

NICMOS Imaging of Molecular Hydrogen Emission in Seyfert Galaxies

A. C. Quillen^{1,2}, & A. Alonso-Herrero¹, M. J. Rieke¹, G. H. Rieke¹, M. Ruiz³, & V. Kulkarni¹

ABSTRACT

We present NICMOS imaging of broad band and molecular hydrogen emission in Seyfert galaxies. In 6/10 Seyferts we detect resolved or extended emission in the 1-0 S(1) λ 2.121 μ m or 1-0 S(3) λ 1.9570 μ m molecular hydrogen lines. We did not detect emission in the most distant galaxy or in the 2 Seyfert 1 galaxies in our sample because of the luminosity of the nuclear point sources. In NGC 5643, NGC 2110 and MKN 1066, molecular hydrogen emission is detected in the extended narrow line region on scales of a few hundred pc from the nucleus. Emission is coincident with [OIII] and H α + [NII] line emission. This emission is also near dust lanes observed in the visible to near-infrared color maps suggesting that a multiphase medium exists near the ionization cones and that the morphology of the line emission is dependent on the density of the ambient media. The high 1-0 S(1) or S(3) H₂ to H α flux ratio suggests that shock excitation of molecular hydrogen (rather than UV fluorescence) is the dominant excitation process in these extended features. In NGC 2992 and NGC 3227 the molecular hydrogen emission is from 800 and 100 pc diameter ‘disks’ (respectively) which are not directly associated with [OIII] emission and are near high levels of extinction ($A_V \gtrsim 10$). In NGC 4945 the molecular hydrogen emission appears to be from the edge of a 100 pc superbubble. In these 3 galaxies the molecular gas could be excited by processes associated with local star formation. We confirm previous spectroscopic studies finding that no single mechanism is likely to be responsible for the molecular hydrogen excitation in Seyfert galaxies.

Subject headings: galaxies: Seyfert — galaxies: molecular hydrogen —

1. Introduction

The emission from molecular hydrogen (H₂) can be used to probe the distribution of excited dense molecular gas in AGNs (active galactic nuclei). H₂ emission (e.g., the 1-0 S(1) line at 2.121 μ m) is known to be bright in Seyferts (e.g., Veilleux et al. 1997; Ruiz 1997; Mouri 1994;

¹The University of Arizona, Steward Observatory, Tucson, AZ 85721

²E-mail: aquillen@as.arizona.edu

³Department of Physical Sciences, University of Hertfordshire, College Lane, Hatfield, Herts AL10 9AB, UK

Koornneef & Israel 1996). Ultraviolet (UV) fluorescence (e.g., Black & van Dishoeck 1987), excitation by low velocity shocks (e.g., Draine & Roberge 1982; Hollenbach & Shull 1977), and heating caused by X-rays are the primary emission mechanisms considered for the excitation of molecular hydrogen (e.g., Puxley et al. 1990; Veilleux et al. 1997; Mouri 1994, Moorwood & Oliva, and references therein). These mechanisms require a source of dense gas (density $\gtrsim 10^3 \text{ cm}^{-3}$) to be either located near a source of illumination (an AGN or a starburst) or actively affected by slow shocks induced either by jets or by kinetic processes (winds, superwinds and supernovae) resulting from star formation. Ground-based spectroscopic studies of Seyferts and LINERS have found that the 1-0 S(1) flux weakly correlates with [FeII] 1.64 μm emission (e.g., Larkin et al. 1998; Ruiz 1997; Mouri 1994) and 6cm radio flux density (Forbes & Ward 1993; Koornneef & Israel 1996; Veilleux et al. 1997) but with a substantial scatter. No correlation between the 1-0 S(1) line and hard X-ray flux in Seyferts has been observed (Ruiz 1997; Veilleux et al. 1997) suggesting that heating by X-rays cannot be the dominant cause of excitation. These results suggest that more than one process causes the H_2 emission.

High resolution imaging of line emission makes it possible to discriminate between likely emission causes. For example, the morphological association between the narrow line region emission lines (e.g., [OIII] $\lambda 5007\text{\AA}$) gas and the radio emission suggests a connection between radio ejecta and the ambient gas (e.g., Wilson & Willis; Unger et al. 1987; Pogge 1988). However, the difference in angle between jet axes, ionization axes and galaxy axes (Nagar et al. 1999) suggest that the distribution of ambient gas in the galaxy also plays an important role in determining the morphology of the ionization cones (also see Quillen et al. 1999 for examples of direct spatial associations). The distribution of extended molecular emission therefore provides a clue to the source of dense gas either illuminated by the central source, affected by energetic motions caused by a jet, or excited by processes associated with star formation (e.g., Maiolino et al. 1997).

In this paper we present a high angular resolution imaging study of molecular hydrogen emission in Seyfert galaxies. Images were observed with the NICMOS cameras on board the Hubble Space Telescope (HST). We construct high angular resolution visible to infrared color maps from WFPC2 and NICMOS broad band images. Red features caused by extinction in these maps are likely to trace dust lanes containing cold dense gas. We compare the morphology of the molecular hydrogen emission with that seen in the color maps and when possible that of [OIII] and $\text{H}\alpha + [\text{NII}]$ emission from ionized gas.

2. Observations

Our sample consists of 3 Seyfert 1 and 7 Seyfert 2 galaxies at a variety of inclinations. We chose nearby Seyferts with bright 1-0 S(1) H_2 fluxes from Ruiz (1997), Veilleux et al. (1997) and Koornneef & Israel (1996). Redshifts were restricted so that either the 1-0 S(1) or 1-0 S(3) line could be observed with existing NICMOS narrow band filters. The H_2 1-0 S(3) 1.9576 μm line is approximately as bright as the 1-0 S(1) line at 2.1213 μm in Seyfert galaxies (Black & van

Dishoeck 1987). The sample is listed in Table 1.

Because the continuum in these galaxies is bright and contains structure which can vary with position, we required observations with a narrow band filter near (in wavelength) to the narrow band filter containing the emission line to accurately subtract the continuum. One HST orbit was spent per galaxy with approximately 128s spent in the broad band filter (F205W or F222M) when possible and the remainder of the orbit evenly divided between the two narrow band filters.

The galaxies were observed in each filter at 4 different positions on the sky with a separation of $0''.57$ or $1''.12$ apart for Camera 2 and 3 observations respectively. Images were reduced with the nircd data reduction software (McLeod 1997) using on orbit darks and flats. Each set of 4 images in a given filter was then combined according to the position observed. The pixel size for the NICMOS camera 2 is $\sim 0''.076$ and for the NICMOS camera 3 is $\sim 0''.204$. Flux calibration for the NICMOS images was performed using the conversion factors based on measurements of the standard stars P330-E and P172-D during the Servicing Mission Observatory Verification program and subsequent observations (M. Rieke, private communication).

2.1. Broad band and archival images

For most of the galaxies we were able to obtain broad band near-IR NICMOS images. Three of the galaxies (MKN 6, NGC 5643, and MKN 1066) were part of snap shot observation programs to observe at $1.60\mu\text{m}$ (in the F160W filter). NGC3227 and IC4329A are part of a GTO (guaranteed time observing) program. For comparison to the F160W broad band images we were able to include as part of our program observations at $2.05\mu\text{m}$ (in the F205W filter) of MKN 1066, NGC 4945, NGC 5506 and NGC 5643 and at $2.22\mu\text{m}$ (in the F222M filter) of MKN 938.

Broad band WFPC2 images in F814W (at $0.814\mu\text{m}$ for NGC 5643) and F606W ($0.606\mu\text{m}$ for NGC 2110, NGC 5643, NGC 2992, MKN 938, MKN 1066 and MKN 6) were retrieved from the HST archive and used with NICMOS images to create optical to infrared color maps. When possible archive images were used to display structure in the ionization cones in either $\text{H}\alpha + [\text{NII}]\lambda 6548$, $6583\text{\AA}$ or $[\text{OIII}]\lambda 5007\text{\AA}$. For the studies discussing these HST data see Mulchaey et al. (1994) on NGC 2110, Simpson et al. (1997) on NGC 5643, Bower et al. (1995) on MKN 1066, and Capetti et al. (1995) on MKN 6.

2.2. Narrow band images

The molecular hydrogen emission was observed with the F212N, F215N, F216N filters on Camera 2 or the F196N on Camera 3 depending on the observed (redshifted) wavelength of the object (see Table 1). For some of the objects in our list, the 1-0 S(1) $2.1213\mu\text{m}$ line could not be observed but the 1-0 S(3) $1.9570\mu\text{m}$ line which is nearly as bright as the 1-0 S(1) line could

be observed with the F196N filter on Camera 3. For all of the objects observed with the F196N filter the line is on the red side of the filter so there is no danger of confusion caused by the [SiVI] $1.96\mu\text{m}$ line which would be outside the filter and should be narrow in the Seyfert 2s (Giannuzzo, Rieke & Rieke 1995). NGC 3227 could be observed in either H_2 line but it has broad lines. To reduce possible confusion from the [SiVI] we chose to observe the 1-0 S(1) line. For this object the line lies at a position in the filter with only 60% of maximum transmission but the object is bright enough so that good signal to noise was achieved with the same integration time as the other objects.

To construct H_2 line emission maps, the narrow band continuum image (lacking the line) was scaled and subtracted from the narrow band image containing the line. From broad band images we estimated the spectral index (or color) of the continuum. We then checked that variations in this index were unlikely to be responsible for the variations seen in the continuum subtracted image. This was the case for all galaxies except for NGC 2992 and except for the nuclei where the structure of the point spread function makes it difficult to measure intensities and colors. In NGC 2992 about 50% of the variations seen in the continuum subtracted image could have been a result of large changes in the continuum spectral index due to reddening. We therefore corrected the continuum image before subtracting it to account for variations in this index. The factor as a function of position used to correct each pixel in this image was derived from the spectral index which we measure from the ratio of the F606W and F205W images of this galaxy. In no cases do the resulting line emission maps show structure observed in the color maps. This suggests that they do indeed trace the distribution of the 1-0 S(1) or S(3) lines. To estimate the line fluxes we took into account the transmission of the filter at the redshifted wavelength of the line. Fluxes and surface brightnesses were measured in various regions from the continuum subtracted images and are listed in Table 2. Fluxes agree within a factor of 2 of the published values (listed in Table 1) in similar apertures. We measure higher nuclear fluxes for NGC 2992 and NGC 5643 than the listed values. Some of the discrepancy could be due to extreme nuclear continuum colors.

2.2.1. NGC 3227

NGC 3227 contains a bright nuclear point source which hampers our ability to detect extended line emission. We attempted to subtract a nuclear point source using a model point spread function (PSF) created by Tiny Tim (Krist & Hook 1997, Krist et al. 1998). However the point source subtracted narrow band images did not produce a line emission map significantly better than that using the images containing the nuclear point sources. Molecular hydrogen emission elongated about $\text{PA} \sim 100^\circ$ is detected in both PSF subtracted and non subtracted line emission maps. The most straightforward interpretation for the extended H_2 emission is that it lies in a 100 pc diameter ‘disk’ with a major axis $\text{PA} \sim 100^\circ$. This ‘disk’ is not aligned with the ionization cone observed in [OIII] at $\text{PA} \sim 30^\circ$ or that of the 18cm radio emission at $\text{PA} \sim -10^\circ$ (Mundell et al. 1995). The position angle of the H_2 emission also differs from that of the major axis of the

galaxy ($\text{PA} \sim 155^\circ$). There is evidence from the CO velocity field that a warped molecular disk is present in this region (Schinnerer et al. 1999). The color map shown in Fig. 1 suggests that extinction is large along the axis of the molecular hydrogen emission.

2.2.2. NGC 2992

H_2 emission is extended along a $\text{PA} \sim 20^\circ$ similar to that of the major axis of the galaxy. The most straightforward interpretation is that it lies in a 800 pc diameter disk in the plane of the galaxy. $\text{H}\alpha + [\text{NII}]$ emission is primarily detected above and below the plane of the galaxy (Wehrle & Morris 1988).

2.2.3. NGC 5643

Strong H_2 emission is observed near the nucleus as well as extended emission coincident with some of the features seen in line emission such as $[\text{OIII}]$. The overall shape of the line emission is similar to but not coincident with dust lanes seen in the optical/near-infrared (F814W/F160W) color map (Quillen et al. 1999). East of the nucleus the radio jet ($\text{PA} = 87^\circ$) lies along the southern edge of the ionization cone (Simpson et al. 1997).

2.2.4. NGC 2110

The nucleus appears to be quite red in our visible/near-infrared color map (see Fig. 4) and so a straight forward subtraction of the F200N image from the F196N image results in over subtraction of the nucleus. In the 1-0 S(3) line emission map we detect extended features near the nucleus at about $1''$ from the nucleus and in a large loop about $3''$ from the nucleus similar in position and shape to that observed in the $\text{H}\alpha + [\text{NII}]$ and $[\text{OIII}]$ line emission maps (which are discussed in Mulchaey et al. 1994). The overall shape of the line emission is similar to but not coincident with features seen in the optical/near-infrared (F814W/F160W) color map (Quillen et al. 1999) and avoids regions of emission from the S-shaped radio jet at $\text{PA} \sim 10^\circ$ (Mulchaey et al. 1994).

2.2.5. MKN 1066

MKN 1066 has narrow line emission (e.g., $[\text{OIII}]$) in two narrow features that are aligned with radio emission from linear jets at $\text{PA} = 134^\circ$ (Bower et al. 1995). Molecular hydrogen emission is detected in the same region as the brightest $[\text{OIII}]$ emission but appears to be somewhat more extended in the direction perpendicular to the radio jets.

2.2.6. NGC 4945

The lack of [OIII] emission and low $3\mu\text{m}$ (L band) flux density suggest that a starbursting disk (rather than a low luminosity AGN) is responsible for the conical shaped cavity observed in the visible and near-IR images of NGC 4945 (Moorwood et al. 1996). The cavity is interpreted to be a region where a bipolar superwind or bubble has evacuated denser material. Our H_2 image resembles that Moorwood et al. (1996), though more structure is seen. The morphology suggests that it forms a boundary between denser gas and a 100 pc scale ‘bubble’ rather than a cone. Extinction on the southern side of the galaxy is so high at $2\mu\text{m}$ that we would not expect to see molecular hydrogen emission from a southern bubble or cone if it exists.

3. Extended emission

We observed extended H_2 emission from the 1-0 S(1) or S(3) line in 6 Seyfert galaxies: NGC 5643, NGC 3227, NGC 2992, NGC 2110, MKN 1066 and NGC 4945. We do not resolve extended emission in the other galaxies. In MKN 938 this might be because of its distance. For MKN 6, NGC 5506 and NGC 5506, the nuclear sources are so bright that it is difficult to observe any extended structure. Future attempts at modeling the point spread function might yield detections of molecular hydrogen emission, particularly in MKN 6 which has extended [OIII] on a scale that we could resolve if the nuclear source could be subtracted reliably.

3.1. H_2 emission in a dense ‘disk’

In both NGC 3227 and NGC 2992 molecular hydrogen emission is coincident with extremely red continuum colors suggesting that it is associated with large quantities of dense molecular material. The morphology of the H_2 emission in both cases is most easily interpreted as emission from a ‘disk’ of 100 and 800 pc diameter in NGC 3227 and in NGC 2992 respectively. In both galaxies line emission from ionized gas and 6cm radio emission (Wehrle & Morris 1988, Mundell et al. 1995) are observed along a different axis than the H_2 emission. Because this emission is associated with high levels of extinction, and not clearly associated with either the ionization cones or radio jets, it is unlikely that shocks or ionization associated with the jets and ionization cones can directly cause the excitation.

In NGC 3227 one possible interpretation is that the molecular hydrogen emission originates from the edge of a circumnuclear disk which has been observed in carbon monoxide submillimeter interferometry (Schinnerer et al. 1999). However in NGC 3327, variations in the distribution and velocity components observed in [OIII] and $\text{H}\alpha$ + [NII] suggest that star formation is vigorous in its central regions (Gonzalez Delgado & Perez 1997) and so that processes associated with star formation (hot stars and supernovae, e.g., Davies, Sugai, & Ward 1998) could also provide the H_2

excitation.

NGC 2992 also has evidence for active star formation. The galaxy has large scale outflows observed in $H\alpha$ and soft X-rays which either could be driven by jets or by a starburst (Colbert et al. 1998). $H\alpha$ emission is observed hundreds of pcs from the nucleus of the galaxy possibly along spiral arms (e.g., Wehrle & Morris 1988). The extent and narrow scale height of the H_2 emission makes it similar to that observed in the prototypical starburst galaxy M82 (Alonso-Herrero et al. 1999).

3.2. The H_2 to $H\alpha$ ratio

For NGC 2110, NGC 5643, and MKN 1066, H_2 emission is associated with features observed in $H\alpha$ + $[NII]$ line emission maps. We can use the strength of the observed $H\alpha$ emission to estimate the local radiation flux of ionizing photons. If the H_2 emission results from UV fluorescence then this flux is directly related (via a spectral index in the UV) to the flux of UV photons that can disassociate molecular hydrogen (in the range 912-1100Å). Models (e.g., Black & van Dishoeck 1987; Puxley et al. 1990) predict 1-0 S(1) molecular hydrogen line strengths as a function of the incident UV radiation field and the density of the molecular clouds. This results in an $H\alpha$ to 1-0 S(1) line flux ratio typical of HII regions (or as used in Puxley et al. 1990 a $Br\gamma$ to 1-0 S(1) ratio). Here we follow the approach outlined in Davies et al. (1998) to determine if UV fluorescence is a feasible H_2 excitation mechanism.

Using the surface brightness in the $H\alpha$ line ($\sim 10^{-14}$ erg s $^{-1}$ cm $^{-2}$ arcsec $^{-2}$ along the northern loop 3'' from the nucleus from Mulchaey et al. 1994 in NGC 2110, $\sim 5 \times 10^{-15}$ erg s $^{-1}$ cm $^{-2}$ arcsec $^{-2}$ in the diffuse emission along the dust lane from Simpson et al. 1997 in NGC 5643 and 10^{-13} erg s $^{-1}$ cm $^{-2}$ arcsec $^{-2}$ on the north-west side of MKN 1066 from Bower et al. 1995), setting the Balmer recombination rate equal to the number of ionizing photons (as in Mulchaey et al. 1994) and using a UV spectral index of 1.6 (as in Davies et al. 1998) we estimate a UV radiation flux between 912 and 1100Å of 50, 22, and 400 times (for NGC 2110, NGC 5643 and MKN 1066 respectively) the solar neighborhood value of 6×10^{-4} erg s $^{-1}$ cm $^{-2}$. From this and using the models of Black & van Dishoeck (1987) we would predict 1-0 S(1) line fluxes of 7×10^{-17} for NGC 2110, 5×10^{-17} for NGC 5643 and 2×10^{-16} erg s $^{-1}$ cm $^{-2}$ arcsec $^{-2}$ for MKN 1066 (assuming $n_H = 10^4$ cm $^{-3}$). This corresponds to the ratio of the 1-0 S(1) to $H\alpha$ flux density ratio of $\sim 10^{-2}$. This is equivalent to a 1-0 S(1)/ $Br\gamma$ flux ratio ~ 1 and agrees with the higher density models of Puxley et al. (1990). From our H_2 emission maps we estimate the surface brightness to be 10, 6 and 30 times higher than that predicted above for UV fluorescence in NGC 2110, NGC 5643 and MKN 1066 respectively. While extinction of $H\alpha$ may to some extent reconcile the flux estimated for UV fluorescent excitation with that observed we suspect that UV fluorescence cannot be the dominant H_2 excitation process in any of these regions. H_2 excitation by slow shocks is then a more likely process (e.g., Koornneef & Israel 1996; Veilleux et al. 1997; Ruiz 1997).

3.3. H₂ Emission in the extended narrow line region

In NGC 2110, NGC 5643 and MKN 1066 we detect emission from molecular hydrogen which appears to be associated and coincident with features observed in [OIII] and H α + [NII] line emission maps tracing ionized gas. The line widths H₂, while smaller than the widths of [FeII] (Ruiz 1997; Veilleux et al. 1997) are still significantly higher than that expected for the low velocity shocks capable of heating the molecular hydrogen without disassociating the molecules ($\lesssim 50$ km/s). If the H₂ emission results from slow shocks then these shocks may be produced by collisions in a multiphase interstellar medium or in other words a medium with patches of molecular material embedded in a more diffuse atomic medium (e.g., Heckman et al. 1986). The association of emission from ionized gas with emission from molecular hydrogen implies that a multiphase medium must exist in or near these cones.

In NGC 5643 and NGC 2110 the line emission is also near dust lanes observed in optical/near-infrared color maps (Quillen et al. 1999). We see that H₂ emission is coincident with H α + [NII] and [OIII] emission, however the dust lanes seen in the color maps, though similar in shape, appear to be offset from the line emission. If the morphology and orientation of the cone is influenced by the density distribution of the ambient media (e.g., as simulated by Mulchaey, Wilson & Tsvetanov 1996), then we would expect that the line emission would peak at a position on the sky that is somewhat different than the maximum extinction or gas density. In fact the proximity of dust lanes with the line emission is good evidence for a dependence of the line excitation on the density of the ambient medium. We might hope that any future set of models for the excitation processes occurring in the ionization cone might be constrained by the observed association and offsets.

Since the UV radiation field implied by the H α + [NII] and [OIII] emission is relatively intense we expect that molecular gas should be disassociated relatively quickly. Either the molecular material exists outside the UV radiation field, is self-shielded or it must be replenished on timescales of $\sim 10^6 - 10^7$ years. The proximity of molecular material also suggests that that dust is present in the ionization cone. This possibility could be tested with future high resolution mid-IR imaging.

4. Correlations between H₂ emission and radio and hard X-ray flux

By comparing the H₂ flux with that in hard X-rays and 6cm radio, previous spectroscopic studies (Veilleux et al. 1997, Koornneef & Israel 1996, Vanzi, Alonso-Herrero & Rieke 1998 and Ruiz 1997) have concluded that more than one emission mechanism is likely to cause the molecular hydrogen emission in Seyfert galaxies. Our imaging study has shown that this is indeed likely. However these studies were hampered by small samples so we take the opportunity here to combine the samples of these previous studies. Molecular hydrogen measurements from Ruiz (1997) are listed in Table 3.

In Fig. 11 we plot 6cm and hard X-ray fluxes vs molecular hydrogen fluxes. A weak correlation noted previously (Ruiz 1997, Forbes & Ward 1993, Koornneef & Israel 1996) between 6cm and molecular hydrogen emission is seen. We note that the scatter is much larger than that observed in [FeII] vs 6cm flux (e.g., Forbes & Ward 1993; Simpson et al. 1996). For most galaxies the hard X-ray flux can heat the molecular gas sufficiently to result in the observed molecular hydrogen emission, however the lack of correlation between the hard X-ray and molecular hydrogen rules out this mechanism as the dominant one.

The weak correlation between H_2 and 6cm emission and lack of correlation with the hard X-ray flux suggests that more than one process causes the molecular hydrogen emission in Seyfert galaxies. Our imaging study suggests that in some of these galaxies a substantial fraction of the emission could be associated with star formation. Unfortunately in starbursting galaxies the flux of H_2 emission does not correlate with any other indicator suggesting that processes other than supernovae and young stars are important (C. Engelbracht, private communication; Vanzi et al. 1998). To test this we looked for (and failed to find) any correlation between the IRAS 12-25 μ m spectral index (which should be dependent on fraction of infrared emission from the nucleus) and the ratio of H_2 emission and hard X-ray flux and the ratio of H_2 and 6cm fluxes.

5. Summary and Discussion

In this paper we have presented NICMOS/HST imaging of molecular H_2 in the 1-0 S(1) or 1-0 S(3) emission in Seyfert galaxies detecting extended emission in 6/10 galaxies. We did not detect extended emission in MKN 938 possibly because of its distance and NGC 5506, IC 4329A and MKN 6 probably because of the luminosity of their nuclear sources. Improvements in techniques for subtraction of NICMOS point spread function (e.g., Krist et al. 1998) may make it possible to detect extended molecular hydrogen emission from these images, particularly in MKN 6.

In NGC 2992 and NGC 3227 the molecular hydrogen emission is from 800 and 100 pc diameter ‘disks’ (respectively) which are not coincident with [OIII] or $H\alpha$ emission and are near high levels of extinction. In NGC 4945 the molecular hydrogen emission appears to be from the edge of a 100 pc superbubble. In these 3 galaxies the molecular gas could be excited by processes associated with local star formation. In NGC 3227 the emission might arise from the edge of a circumnuclear torus or warped disk.

In NGC 5643, NGC 2110 and MKN 1066, molecular hydrogen emission is detected in the extended narrow line region on scales of a few hundred pc from the nucleus. Emission is found in the ionization cones coincident with [OIII] and $H\alpha$ + [NII] line emission. This emission is also near dust lanes observed in the visible to near-infrared color maps. The coincidence of H_2 emission with that from ionized gas implies that a multiphase medium is near or within the ionization cone. The proximity to dust lanes seen in the color maps suggests that processes causing the line emission are dependent on the density of the ambient media. The high 1-0 S(1) or S(3) H_2 to

H α flux ratio suggests that shock excitation (not UV fluorescence) of molecular hydrogen is the dominant excitation process in these extended features.

We have compiled the spectroscopic observations of molecular hydrogen (from Veilleux et al. 1997, Koornneef & Israel 1996 and Ruiz 1997) to look for correlations between 6cm radio and hard X-ray flux. We find no correlation between H₂ emission and hard X-ray flux confirming the results of these previous studies, and ruling out heating by X-rays as the dominant source of H₂ excitation. There is only a weak correlation between H₂ emission flux and 6cm flux density. Both our imaging study and this compilation confirm the finding of previous spectroscopic studies (Ruiz 1997, Koornneef & Israel 1996, Veilleux et al. 1997, Forbes & Ward 1993): no single mechanism is likely to be responsible for the molecular hydrogen excitation in Seyfert galaxies. Future higher resolution observations may allow us to separate between likely processes and isolate molecular hydrogen emission from the circumnuclear environment.

Support for this work was provided by NASA through grant number GO-07869.01-96A from the Space Telescope Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. We also acknowledge support from NASA project NAG-53359. We acknowledge helpful discussions and correspondence with A. Eckart, E. Schinnerer, C. Kulesa, G. Bower and A. Cotera.

REFERENCES

- Alonso-Herrero, A., Rieke, M. J., et al. 1999, in preparation
- Bassani, L., Dadina, M., Maiolino, R., Salvati, M. Risaliti, G., Della Ceca, R., Matt, G., & Zamorani, G. 1999, ApJS, in press, (astro-ph/9811074)
- Black, J. H., & van Dishoeck, E. F. 1987, ApJ, 322, 412
- Bower, G., Wilson, A., Morse, J. A., Gelderman, R., Whittle, M., & Mulchaey, J. 1995, ApJ, 454, 106
- Capetti, A., Axon, D. J., Kukula, M., Macchetto, F., Pedlar, A., Sparks, W. B., & Boksenberg, A. 1995, ApJ, 454, L85
- Colbert, E. J. M., Baum, S. A., O’Dea, C. P., & Veilleux, S. 1998, ApJ, 496, 786
- Davies, R. I., Sugai, H., & Ward, M. J. 1998, MNRAS, 300, 388
- Draine, B. T., & Roberge, W. G. 1982, ApJ, 259, L91
- Forbes, D. A., & Ward, M. J. 1993, ApJ, 416, 150
- Giannuzzo, E., Rieke, G. H., & Rieke, M. J. 1995, ApJL, 446, L5
- Gonzalez Delgado, R. M., & Perez, E. 1997, MNRAS, 284, 931
- Heckman, T. M., Beckwith, S., Blitz, L., Skrutskie, M., & Wilson, A. S. 1986, ApJ, 305, 157
- Hollenbach, D. J., & Shull, J. M. 1977, ApJ, 216, 419
- Koornneef, J., & Israel, F. P. 1996, New Astro., 1, 271
- Krist, J., & Hook, R. 1997, ‘The Tiny Tim User’s Guide’, <http://scivax.stsci.edu/~krist.tinytim.html>
- Krist, J. E., Golimowski, D. A., Schroeder, D. J., & Henry, T. J. 1998, PASP, 110, 1046
- Larkin, J. E., Armus, L., Knop, R. A., Soifer, B. T., & Matthews, K. 1998, ApJS, 114, 59
- Lepp, S., & McCray, R. 1983, ApJ, 269, 560
- Maiolino, R., Ruiz, M., Rieke, G. H., & Papadopoulos, P. 1997, ApJ, 485, 552
- McLeod, B. 1997, proceedings of the 1997 HST Calibration Workshop, eds. S. Casertano, R. Jędrzejewski, T. Keyes, and M. Stevens, published by the Space Telescope Science Institute, Baltimore, MD, p. 281
- Moorwood, A. F. M., van der Werf, P. P., Kotilainen, J. K., Marconi, A., & Oliva, E. 1996, A&A, L1

- Moorwood, A. F. M., & Oliva, E. 1988, *A&A*, 203, 278
- Mouri, H. 1994, *ApJ*, 427, 777
- Mulchaey, J. S., Wilson, A. S., Bower, G. A., Heckman, T. M., Krolik, J. H., & Miley, G. K. 1994, *ApJ*, 433, 625
- Mulchaey, J. S., Wilson, A. S., & Tsvetanov, Z. 1996, *ApJ*, 467, 197
- Mundell, C. G., Holloway, A. J., Pedlar, A., Meaburn, J., Kukula, M. J., & Axon, D. J. 1995, *MNRAS*, 275, 67
- Nagar, N. M., Wilson, A. S., Mulchaey, J. S., & Gallimore, J. F. 1999, *ApJS*, in press
- Pogge, R. W. 1988, *ApJ*, 332, 702
- Puxley, P. J., Hawarden, T. G., & Mountain, C. M. 1990, *ApJ*, 364, 77
- Quillen, A. C., Alonso-Herrero, A., Rieke, M. J., McDonald, C., Falcke, H., & Rieke, G. H. 1999, *ApJ*, in press, (astro-ph/9902233)
- Reynolds, C. S. 1997, *MNRAS*, 286, 513
- Ruiz, M. 1997, PhD Thesis, University of Arizona
- Schinnerer, E., Eckart, A., & Tacconi, L. J. 1999, ‘Galaxy Dynamics’, proceedings of a conference held at Rutgers University, 8-12 Aug 1998. To appear in ASP Conference Series, edited by D. R. Merritt, M. Valluri, and J. A. Sellwood
- Simpson, C., Wilson, A. S., Bower, G. Heckman, T. M., Krolik, J. H., & Miley, G. K. 1997, *ApJ*, 474, 121
- Simpson, C., Forbes, D. A., Baker, A. C., & Ward, M. J. 1996, *MNRAS*, 283, 777
- Unger, S. W., Pedlar, A., Axon, D. J., Whittle, M., Meurs, E. J. A., & Ward, M. J. 1987, *MNRAS*, 228, 671
- Vanzi, L., Alonso-Herrero, A., & Rieke, G. H. 1998, *ApJ*, 504, 93
- Veilleux, S., Goodrich, R. W., & Hill, G. J. 1997, *ApJ*, 477, 631
- Wehrle, A. E., & Morris, M. 1988, *AJ*, 95, 1689
- Wilson, A. S., & Willis, A. G. 1980, *ApJ*, 240, 429

Fig. 1.— a) NICMOS Camera 2 F160W ($1.60\mu\text{m}$) image of NGC 3227. Contours 0.25 mag apart are overlayed with the faintest white counter roughly equivalent to $14.0 \text{ mag}/''^2$ in Johnson H band. b) Visible/near-IR color map constructed from F606W and F160W images. Colors range from $V-H=2.6$ (approximately Johnson bands; white) to $V-H=4.1$ (black). c) F606W ($0.606\mu\text{m}$) WFPC2 image. Contours are 0.25 mag apart. Note that the nucleus is saturated in this image. d) 1-0 S(1) H_2 line emission. This image was constructed using PSF subtracted images. Dark colors refer to brighter emission. $1''$ is equivalent to 75 pc assuming $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$. North is up and east is to the left.

Fig. 2.— a) NICMOS Camera 2 F205W ($2.05\mu\text{m}$) image of NGC 2992. Contours 0.25 mag apart are overlayed with the faintest white counter roughly equivalent to $4.61 \text{ mJy}/''^2$. b) Visible/near-IR color map constructed from F606W and F205W images. Colors range from $V-K=4$ (approximately Johnson bands; white) to $V-K=7$ (black). c) F606W ($0.606\mu\text{m}$) WFPC2 image. Contours are 0.25 mag apart. Note that the nucleus is saturated in this image. d) 1-0 S(3) H_2 line emission constructed from NICMOS Camera 3 narrow band images. Dark colors refer to brighter emission. $1''$ is equivalent to 153 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 3.— a) NICMOS Camera 2 F160W ($1.60\mu\text{m}$) image of NGC 5643. Contours 0.25 magnitude apart are overlayed. The faintest white contour corresponds to $14.0 \text{ mag}/''^2$ in Johnson H band. b) Visible/near-IR color map constructed from F814W and F160W images. Colors range from $I-H=1.3$ (approximately Johnson bands; white) to $I-H=2.3$ (black). c) $\text{H}\alpha + [\text{NII}]$ emission (from WFPC2 images; see Simpson et al. 1997). d) 1-0 S(3) H_2 line emission map constructed from NICMOS Camera 3 narrow band images. Dark colors refer to brighter emission. $1''$ is equivalent to 78 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 4.— a) NICMOS Camera 3 F200N ($2.00\mu\text{m}$) image of NGC 2110. Contours 0.25 magnitude apart are overlayed with the faintest white counter roughly equivalent to $1.9 \text{ mJy}/''^2$. b) Visible/near-IR color map constructed from F606W and F200N images. Colors range from $V-K=4$ (approximately Johnson bands; white) to $V-K=5.4$ (black). c) $\text{H}\alpha + [\text{NII}]$ emission (from PC images; see Mulchaey et al. 1994). d) 1-0 S(3) H_2 line emission constructed from NICMOS Camera 3 narrow band images. Dark colors refer to brighter emission. $1''$ is equivalent to 148 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 5.— a) NICMOS Camera 2 F160W ($1.60\mu\text{m}$) image of MKN 1066. Contours 0.5 magnitude apart are overlayed with the faintest white contour roughly equivalent to $14.0 \text{ mag}/''^2$ in Johnson H band. b) Visible/near-IR color map constructed from F606W and F160W images. Colors range from $V-H=2.5$ (approximately Johnson bands; white) to $V-H=4.5$ (black). Note that the F606W image is contaminated with emission from $H\alpha$. Light corresponds to blue colors or regions of bright $H\alpha$ emission and dark corresponds to red colors. c) [OIII] line emission (from PC images; see Bower et al. 1995). d) 1-0 S(1) H_2 line emission constructed from NICMOS Camera 2 narrow band images. Dark colors refer to brighter emission. $1''$ is equivalent to 238 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 6.— a) NICMOS Camera 2 F205W ($2.05\mu\text{m}$) image of NGC 4945. Contours 0.25 mag apart are overlayed with the faintest white counter roughly equivalent to $4.61 \text{ mJy}/''^2$. b) 1-0 S(1) H_2 line emission constructed from NICMOS Camera 2 narrow band images. Dark colors refer to brighter emission. $1''$ is equivalent to 36.3 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 7.— a) NICMOS Camera 2 F205W ($2.05\mu\text{m}$ image) of MKN 938. Contours 0.25 magnitude apart are overlayed with the faintest white contour corresponding to $4.21 \text{ mJy}/''^2$. b) Visible/near-IR color map constructed from F606W and F205W images. Colors range from $V-K=2.8$ (approximately Johnson bands; white) to $V-K=4.8$ (black). c) F606W ($0.606\mu\text{m}$) WFPC2 image. d) 1-0 S(1) H_2 line emission constructed from NICMOS Camera 2 narrow band images. We do not detect any extended emission in molecular hydrogen. Dark colors would refer to brighter emission. $1''$ is equivalent to 384 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 8.— a) NICMOS Camera 2 F160W ($1.60\mu\text{m}$) image of NGC 5506. b) Continuum subtracted narrow band image of the 1-0 S(3) H_2 line constructed from NICMOS Camera 3 narrow band images. The images are dominated by the Seyfert 1 point source and we detect no extended line emission. $1''$ is equivalent to 120 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 9.— a) NICMOS Camera 2 narrow band continuum F212N ($2.12\mu\text{m}$) image of IC 4329A. b) Continuum subtracted narrow band image of the 1-0 S(1) H_2 line. The images are dominated by the Seyfert 1 point source and we detect no extended line emission. $1''$ is equivalent to 311 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 10.— a) NICMOS Camera 1 F160W ($1.60\mu\text{m}$) image of MKN 6. b) F606W ($0.606\mu\text{m}$) WFPC2 image. We have displayed this image to show the dust lane to the north. The nucleus is saturated in this image. c) [OIII] line emission (from FOC images; see Capetti et al. 1995). d) Continuum subtracted narrow band image of the 1-0 S(1) H_2 line. The images are dominated by the Seyfert 1 point source and we detect no extended line emission. $1''$ is equivalent to 358 pc (for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Fig. 11.— a) Molecular hydrogen line flux vs 6 cm radio emission. Seyfert 1s are shown as solid triangles and Seyfert 2s as open triangles. We see only a weak correlation between 6cm and molecular hydrogen flux. The weakness of the correlation is similar to that presented in Veilleux et al. (1997) and Ruiz (1997). We note that a correlation was also reported by Forbes & Ward (1993) and Koornneef & Israel (1996). H_2 and 6cm fluxes are compiled from Koornneef & Israel (1996), Ruiz (1997), Veilleux et al. (1997), and Moorwood & Oliva (1988). b) Molecular hydrogen line flux vs hard X-ray flux. Points below and to the right of the solid line have sufficient X-ray flux to account for the heating of the molecular gas (as estimated by Lepp & McCray 1983, $F(HX) = 400f(H_2\lambda 2.121)$; this plot is similar to that shown in Veilleux et al. 1997). However the lack of correlation between molecular hydrogen emission and X-ray flux suggests that heating by X-rays is unlikely to be the dominant heat source for the molecular hydrogen emission. For the Seyfert 2s hard X-ray fluxes are compiled from Bassani et al. (1999) and for the Seyfert 1s from Reynolds (1997).

Table 1. Sample

Galaxy (1)	Inclination (2)	Type (3)	Flux H ₂ (4)	z (5)	Obs. wave (6)	Camera (7)	Line (8)	Continuum (9)
NGC 3227	45	Sy1.5	29.0	0.00386	2.129	NIC2	F212N	F215N
NGC 2992	70	Sy2	12	0.0079	1.973	NIC3	F196N	F200N
NGC 2110	40	Sy2	9.25	0.00762	1.972	NIC3	F196N	F200N
NGC 5643	30	Sy2	21	0.00400	1.965	NIC3	F196N	F200N
NGC 4945	80	Sy2	129	0.00187	2.122	NIC2	F212N	F215N
Mkn 1066	55	Sy2	13.6	0.0123	2.147	NIC2	F215N	F212N
Mkn 938	70	Sy2	17.3	0.01978	2.163	NIC2	F216N	F212N
Mkn 6	50	Sy1.5	19.5	0.01847	2.161	NIC2	F216N	F212N
IC 4329A	75	Sy 1	11.8	0.01605	2.155	NIC2	F215N	F212N
NGC 5506	70	Sy1.9	10.6	0.00618	1.970	NIC3	F196N	F200N

(1) Galaxy; (2) Galaxy inclination in degrees where 0 = face-on; (3) Seyfert type; (4) 1-0 S(1) 2.1213 μ m line flux in units of 10^{-15} erg cm⁻² s⁻¹ compiled in Ruiz (1997), Veilleux et al. (1997), or Koornneef & Israel (1996). All but NGC 5643, Mkn1066, NGC 4945 are from Ruiz (1997) and fluxes were extracted from a 1.2'' slit. NGC 4945 is from Koornneef & Israel (1996) and was from a 6x6'' aperture. MKN 1066 flux is from Veilleux et al. (1997) and was from a 1.5x1.5'' aperture. (5) Redshift; (6) Observable wavelength (redshifted) for the 1-0 S(1) 2.1213 μ m or 1-0 S(3) 1.9576 μ m line; (7) NICMOS camera used for narrow band imaging; (8) Narrow band filter containing the H₂ line. (9) Continuum filter.

Table 2. 1-0 S(1) and 1-0 S(3) H₂ fluxes

Galaxy	Region	Flux	Surface brightness
NGC 5643	eastern feature	1.2'' < r < 8.16''	3
	western feature	1.2'' < r < 8.16''	
	nuclear	r < 1.2''	
NGC 2110	near the nucleus	0.6'' < r < 1.6''	6
	northern loop	2.0 < r < 4.0''	
NGC 2992	nuclear	r < 0.6''	68
	disk	0.6'' < r < 6.5''	
NGC 3227	disk	0.4 < r < 1.3''	10
MKN 1066	total	r < 1.1''	70
	western feature	$r \sim 0.5''$	
NGC 4945	total	170	

NGC 5643, NGC 2110, and NGC 2992 fluxes are for the 1-0 S(3) line. For the remaining galaxies we list fluxes in the 1-0 S(1) line. Fluxes are listed in units of 10^{-15} erg cm⁻² s⁻¹ and surface brightnesses in units of 10^{-16} erg cm⁻² s^{-1''-2}.

Table 3. 1-0S(1)H₂ fluxes and FWHM from Ruiz (1997).

Galaxy	flux 10 ⁻¹⁴ erg cm ⁻² s ⁻¹	FWMH km s ⁻¹
Mkn 334	0.91	130
Mkn 335	0.13	102
Mkn 938	1.73	537
Mkn 348	< 0.04	...
Mkn 1040	0.24	185
NGC 1275	3.71	382
Mkn 1095	0.51	381
NGC 2110	0.93	447
MCG-8-11-11	1.19	386
NGC 2273	0.68	251
Mkn 6	1.95	234
Mkn 376	< 0.02	...
Mkn 79	< 0.05	...
Mkn 704	< 0.06	...
NGC 2992	0.41	327
NGC 3227	2.90	331
NGC 3516	< 0.01	...
NGC 4051	0.87	253
NGC 4151	0.20	232
Mkn 766	0.24	214
IC 4329A	1.18 ^a	332
NGC 5506	1.06	315
Mkn 817	< 0.02	...
NGC 7469	0.86	309
Mkn 533	0.48	268

^aDerived from Giannuzzo et al. (1995).

This figure "page1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page6.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page7.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page8.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page9.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page10.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>

This figure "page11.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/9907011v1>